arXiv:1603.04694v2 [math.CA] 16 Mar 2016

ZEROS OF RAMANUJAN TYPE ENTIRE FUNCTIONS

RUIMING ZHANG

Abstract. In this work we establish some polynomials and entire functions have only real zeros. These polynomials generalize q-Laguerre polynomials $L_n^{(\alpha)}(x;q)$, while the entire functions are generalizations of Ramanujan's entire function $A_q(z)$, q-Bessel functions $J_\nu^{(2)}(z;q)$, $J_\nu^{(3)}(z;q)$ and confluent basic hypergeometric series.

1. Introduction

In [12] Ismail and Zhang defined an entire function

$$A_q^{(\alpha)}(a;z)=\sum_{n=0}^{\infty}\frac{(a;q)_n\,q^{\alpha n^2}z^n}{(q;q)_n}, \tag{1.1}$$

where $\alpha>0$, $0<q<1$, and for any positive integer $m\in\mathbb{N}$ and all complex numbers $a,a_1,\dots,a_m\,n\in\mathbb{C}$, [1, 11]

(1.2)

$$(a;q)_\infty=\prod_{k=0}^{\infty}\left(1-aq^k\right),\ (a;q)_n=\frac{(a;q)_\infty}{(aq^n;q)_\infty},\ (a_1,\dots,a_m;q)_n=\prod_{j=1}^{m}(a_j;q)_n.$$

Clearly, $A_q^{(\alpha)}(a;z)$ is a kind of entire function treated in Lemma 14.1.4 of [11], and by the lemma it must have infinitely many zeros if $aq^n\neq 1$ for any nonnegative integers $n\in\mathbb{N}_0$. Furthermore, in [8] Hayman proved an asymptotic expansion for the n-th zero of this class of entire function under the general condition that the parameter q is strictly inside the unit disk $|q|<1$.

We observe that

$$A_q^{(1/2)}\left(q^{-n};z\right)=\sum_{k=0}^{\infty}\frac{(q^{-n};q)_k\,q^{k^2/2}z^k}{(q;q)_k}=(q;q)_n\,S_n\left(zq^{1/2-n};q\right), \tag{1.3}$$

and

$$A_q^{(1)}(0;z)=\sum_{n=0}^{\infty}\frac{q^{n^2}z^n}{(q;q)_n}=A_q(-z),\ A_q^{(1)}(q;z)=\sum_{n=0}^{\infty}q^{n^2}z^n, \tag{1.4}$$

where $A_q(z)$ and $S_n(z;q)$ are the Ramanujan entire function and Stieltjes-Wigert polynomial respectively, see [11], for example. Since $A_q^{(\alpha)}(a;z)$ generalizes both

2000 *Mathematics Subject Classification.* 33D15; 33C10; 33D99; 33C99.

Key words and phrases. Zeros of polynomials; zeros of q-series; Laguerre-Pólya class; Pólya frequence sequence; q-Laguerre polynomials; q-Bessel functions; basic hypergeometric series.

This research is partially supported by National Natural Science Foundation of China, grant No. 11371294 and Northwest A&F University.

$A_q(z)$ and $S_n(z;q)$, and it is well-known that both of them have all real positive zeros. Thus it is natural to ask under what conditions the zeros of the entire function $A_q^{(\alpha)}(a;z)$ are all real. In this work we shall present a partially answer to this question and more.

2. Preliminaries

In the proofs we need the Vitali's theorem [15] and Hurwitz's theorem [2]. For our convenience we list them here. The first is the Vitali's theorem:

Theorem 1. *Let $\{f_n(z)\}$ be a sequence of functions analytic in a domain D and assume that $f_n(z) \to f(z)$ point-wise in D. Then $f_n(z) \to f(z)$ uniformly in any subdomain bounded by a contour C, provided that C is contained in D.*

Here is the Hurwitz's theorm:

Theorem 2. *If the functions $f_n(z)$ are analytic and $\neq 0$ in a region Ω, and if $f_n(z)$ converges to $f(z)$, uniformly on every compact subset of Ω, then $f(z)$ is either identically zero or never equal to zero in Ω.*

It is known that if an entire function has a finite order that it is not a positive integer, then $f(z)$ has infinitely many zeros, [4, 11]. Given an entire function of order 0 with $f(0) \neq 0$, if $-z_k,\ k \in \mathbb{N}$ are all the roots, then by Hadamard's canonical representation of entire functions we have

$$f(z) = f(0) \prod_{k=1}^{\infty} \left(1 + \frac{z}{z_k}\right), \tag{2.1}$$

where $\sum_{k=1}^{\infty} |z_k|^{-1} < \infty$. A real entire function $f(z)$ is of Laguerre-Pólya class if [6, 10]

$$f(z) = cz^m e^{-\alpha z^2 + \beta z} \prod_{k=1}^{\infty} \left(1 + \frac{z}{z_k}\right) e^{-z/z_k}, \tag{2.2}$$

where $c, \beta,\ z_k \in \mathbb{R},\ \alpha \geq 0,\ m \in \mathbb{Z}^+$, and $\sum_{k=1}^{\infty} z_k^{-2} < \infty$. Clearly, given a real entire function $f(z)$ of order 0 that has nonnegative Taylor coefficients and satisfies $f(0) \neq 0$, if it is also in the Laguerre-Pólya class then it must have the factorization (2.1) with $z_k > 0$.

A real sequence $\{a_n\}_{n=0}^{\infty}$ is called a Pólya frequence (PF) sequence if the infinite matrix $(a_{j-i})_{i,j=0}^{\infty}$ is totally positive, i.e. all its minors are nonnegative, where we follow the usual convention that $a_n = 0$ if $n < 0$. [3, 7, 10]

Theorem 3. *The sequence $\{b_k\}_{k=0}^{\infty}$ is a PF sequence if and only if*

$$cz^m e^{\gamma z} \prod_{k=1}^{\infty} \frac{1 + \alpha_k z}{1 - \beta_k z} = \sum_{k=0}^{\infty} b_k z^k, \tag{2.3}$$

where $c \geq 0,\ \gamma \geq 0,\ \alpha_k \geq 0,\ \beta_k \geq 0$, $m \in \mathbb{Z}^+$, and $\sum_{k=1}^{\infty} (\alpha_k + \beta_k) < \infty$.

In the case $b_k = 0$ for $k = n+1, \ldots$, then the right hand series becomes a polynomial of degree at most n, then we have the following, [3, 7, 10, 14]

Theorem 4. *The PF sequences have the following properties:*

(1) *Let $b_k \geq 0,\ k \geq 0$. Then, the sequence $\{b_k\}_{k=0}^{n}$ is a PF sequence if and only if the polynomial $\sum_{k=0}^{n} b_k x^k$ has all nonpositive zeros.*
(2) *Let $a_k,\ b_k \geq 0,\ k \geq 0$. If $\{a_k\}_{k=0}^{m}$ and $\{b_k\}_{k=0}^{n}$ are PF sequences, then so is $\{a_k b_k\}_{k\geq 0}$.*
(3) *Let $b_0,\ b_1,\ \ldots,\ b_n \geq 0$ and $\{b_k\}_{k=0}^{n}$ be a PF sequence, then so is $\{b_k/k!\}_{k=0}^{n}$.*
(4) *Let $\{a_k\}_{k=0}^{m}$ and $\{b_k\}_{k=0}^{n}$ be PF sequences such that $a_k,\ b_k \geq 0,\ k \geq 0$, then $\{k! a_k b_k\}_{k\geq 0}$ is also a PF sequence.*

It was proved in that [5, 6]

$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{n!} x^n \tag{2.4}$$

is in Laguerre-Pólya class for $q \in (-1, 1)$. For $0 < q < 1$. This function is clearly a real entire function of order 0 with positive Taylor coefficients, hence the sequence $\left\{q^{n^2}/n!\right\}_{n=0}^{\infty}$ is a PF sequence by Theorem 3.

For $a \geq 0$ and $0 < q < 1$, from the q-Binomial theorem we have [1, 11]

$$\frac{(-az;q)_\infty}{(bz;q)_\infty} = \sum_{n=0}^{\infty} \frac{(-a;q)_n}{(q;q)_n} z^n, \tag{2.5}$$

and

$$(-q^n z;q)_n = \sum_{k=0}^{n} \frac{(q^{-n};q)_k (-z)^k}{(q;q)_k}. \tag{2.6}$$

Then by Theorem 3 both$\{(-a;q)_k/(q;q)_k\}_{k=0}^{\infty}$ and $\left\{(-1)^k (q^{-n};q)_k /(q;q)_k\right\}_{k=0}^{\infty}$ are PF sequences.

The q-Bessel functions $J_\nu^{(1)}(z;q)$, $J^{(2)}(z;q)$ and $J^{(3)}(z;q)$ are defined by [11]

$$\frac{2^\nu (q;q)_\infty J_\nu^{(1)}(z;q)}{(q^{\nu+1};q)_\infty z^\nu} = \sum_{n=0}^{\infty} \frac{\left(-z^2/4\right)^n}{(q, q^{\nu+1};q)_n}, \tag{2.7}$$

$$\frac{2^\nu (q;q)_\infty J_\nu^{(2)}(z;q)}{(q^{\nu+1};q)_\infty z^\nu} = \sum_{n=0}^{\infty} \frac{q^{n^2}\left(-z^2 q^\nu/4\right)^n}{(q, q^{\nu+1};q)_n}, \tag{2.8}$$

and

$$\frac{2^\nu (q;q)_\infty J_\nu^{(3)}(z;q)}{(q^{\nu+1};q)_\infty z^\nu} = \sum_{n=0}^{\infty} \frac{q^{\binom{n+1}{2}}\left(-z^2/4\right)^n}{(q, q^{\nu+1};q)_n} \tag{2.9}$$

respectively.

For $\nu > -1$ and $0 < q < 1$, it is known that $J^{(2)}(z;q)$ is an entire function of order 0 such that

$$\frac{2^\nu (q;q)_\infty J_\nu^{(2)}(z;q)}{(q^{\nu+1};q)_\infty z^\nu} = \prod_{n=0}^{\infty} \left(1 - \frac{z^2}{j_{\nu,n}(q)^2}\right), \tag{2.10}$$

where

$$0 < j_{\nu,1}(q) < j_{\nu,2}(q) < \dots \tag{2.11}$$

are the positive zeros of $J^{(2)}(z;q)$ satisfying

$$\sum_{n=1}^{\infty} \frac{1}{j_{\nu,n}(q)^2} < \infty. \tag{2.12}$$

For $\nu > -1$ and $0 < q < 1$, from (2.10) and the relation [11]

$$J_\nu^{(1)}(z;q) = \frac{J_\nu^{(2)}(z;q)}{(-z^2/4;q)_\infty} \tag{2.13}$$

we get

$$\sum_{n=0}^{\infty} \frac{z^n}{(q,q^{\nu+1};q)_n} = \frac{\prod_{n=0}^{\infty}\left(1+z/j_{\nu,n}(q)^2\right)}{(z/4;q)_\infty}. \tag{2.14}$$

From (2.8), (2.10) and (2.14) we see that both $\left\{1/\left(q,q^{\nu+1};q\right)_k\right\}_{k=0}^{\infty}$ and $\left\{q^{k^2}/\left(q,q^{\nu+1};q\right)_k\right\}_{k=0}^{\infty}$ are PF sequences by Theorem 3.

It is worth noticing that these q-Bessel functions are much more than merely being q-analogues to their classical counterpart

$$J_\nu(z) = \frac{(z/2)^\nu}{\Gamma(\nu+1)} \sum_{n=0}^{\infty} \frac{\left(-z^2/4\right)^n}{n!(\nu+1)_n}, \tag{2.15}$$

where

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)}, \quad a,n \in \mathbb{Z} \tag{2.16}$$

with $-a-n \notin \mathbb{N}_0$ and $\Gamma(z)$ is the Euler's Gamma function. They have many interesting properties that the classical $J_\nu(z)$ does not possess. Here we only mention two of them. First of them is that $J_\nu^{(2)}\left(2iq^{-n/2};q\right),\ n \in \mathbb{N}_0$ can be evaluated explicitly, see [13]. The second is that there are many symmetries among themselves.

Let

$$j_\nu^{(k)}(z;q) = \frac{2^\nu (q;q)_\infty J_\nu^{(k)}(z;q)}{(q^{\nu+1};q)_\infty\, z^\nu}, \quad k = 1,2,3. \tag{2.17}$$

Then for $q > 1$ and $|z| < 1$, we have

$$j_\nu^{(1)}\left(z;\frac{1}{q}\right) = j_\nu^{(2)}\left(\sqrt{q}z;q\right),\ j_\nu^{(2)}\left(z;\frac{1}{q}\right) = j_\nu^{(1)}\left(\sqrt{q}z;q\right),\ j_2^{(3)}\left(z;\frac{1}{q}\right) = j_\nu^{(3)}\left(q^{\frac{\nu}{2}}z;q\right). \tag{2.18}$$

The above relations are very typical among basic hypergeometric series, but it is seldom explored in the literature.

3. Main Results

We follow the usual convention that an empty sum is zero but an empty product is 1.

Theorem 5. *Let $n \in \mathbb{N}$, $0 < q < 1$, and $\alpha \geq 0$.*

(1) *The polynomial*

$$A_q^{(\alpha)}\left(q^{-n};x\right)=\sum_{k=0}^{n}\frac{\left(q^{-n};q\right)_k q^{\alpha k^2}x^k}{(q;q)_k} \tag{3.1}$$

has all positive zeros.

(2) *For $m,\ell\in\mathbb{N}_0$, $n_j\in\mathbb{N}$, $0<q_j<1$, $1\le j\le m$, and $\beta_r>0$, $0<q_r<1$, $1\le r\le\ell$, the polynomial*

$$\sum_{k=0}^{\min\{n_j|1\le j\le m\}}\prod_{j=1}^{m}\frac{\left(q_j^{-n_j};q_j\right)_k}{(q_j;q_j)_k}\frac{q^{\alpha k^2}\left((-1)^m x\right)^k}{\prod_{r=1}^{\ell}\left(q_r,q_r^{\beta_r};q_r\right)_k} \tag{3.2}$$

has all negative zeros.

(3) *For $m,\ell\in\mathbb{N}_0$, $n_j\in\mathbb{N}$, $1\le j\le m$, and $\beta_r>0$, $1\le r\le\ell$, the polynomial*

$$\sum_{k=0}^{\min\{n_j|1\le j\le m\}}\frac{\prod_{j=1}^{m}(-n_j)_k}{\prod_{r=1}^{\ell}(\beta_r)_k}\frac{\left((-1)^m x\right)^k}{(k!)^{m+\ell}} \tag{3.3}$$

has all negative zeros.

The following results are for the entire functions in the first parameter range:

Theorem 6. *Let $0<q<1$ and $\alpha>0$. Then,*

(1) *For $a\ge 0$, the entire function*

$$A_q^{(\alpha)}\left(-a;z\right)=\sum_{n=0}^{\infty}\frac{\left(-a;q\right)_n q^{\alpha n^2}z^n}{(q;q)_n} \tag{3.4}$$

has infinitely many zeros and all of them are negative.

(2) *For $m,\ell\in\mathbb{N}_0$, $a_j\ge 0$, $0<q_j<1$, $1\le j\le m$, and $\beta_r>0$, $0<q_r<1$, $1\le r\le\ell$, the entire function*

$$\sum_{k=0}^{\infty}\prod_{j=1}^{m}\frac{\left(-a_j;q_j\right)_k}{(q_j;q_j)_k}\frac{q^{\alpha k^2}z^k}{\prod_{r=1}^{\ell}\left(q_r,q_r^{\beta_r};q_r\right)_k} \tag{3.5}$$

has infinitely many zeros and all of them are negative.

(3) *For $m\in\mathbb{N}_0$, $\ell\in\mathbb{N}$ and $\beta_r\ge 1$, $1\le r\le\ell$, the entire function*

$$\sum_{k=0}^{\infty}\frac{z^k}{(k!)^{m+\ell}\prod_{r=1}^{\ell}\left(\beta_r\right)_k} \tag{3.6}$$

has infinitely many zeros and all of them are negative.

For $r,\ s\in\mathbb{N}_0$ and $0<q<1$, let $a_1,\ldots,\ a_r,\ b_1,\ldots,\ b_s\in\mathbb{C}$

$${}_rA_s^{(\alpha)}(a_1,\ldots a_r;\ b_1,\ldots,\ b_s\ ;\ q;\ z)=\sum_{n=0}^{\infty}\frac{(a_1,\ldots,\ a_r;\ q)_n}{(b_1,\ldots,b_s;\ q)_n}q^{\alpha n^2}z^n, \tag{3.7}$$

then it is clear that for $\alpha>0$ and $0<q<1$ we have

$$A_q^{(\alpha)}(a;z)=_1 A_1^{(\alpha)}(a,\ q;z), \tag{3.8}$$

and for $z \in \mathbb{C}$ and $s+1 > r$ we have

$$\begin{aligned} &{}_rA_s^{((s+1-r)/2)}\left(a_1,\dots a_r;\ q,\ b_1,\dots,\ b_s\ ;\ q;\ (-1/\sqrt{q})^{s+1-r}z\right) \\ &= {}_r\phi_s\left(\begin{array}{c} a_1,\dots,\ a_r \\ b_1,\dots,\ b_s \end{array}\middle| q, z\right), \end{aligned} \tag{3.9}$$

where the basic hypergeometric series ${}_r\phi_s$ is defined by [1, 11]

$$ {}_r\phi_s\left(\begin{array}{c} a_1,\dots,\ a_r \\ b_1,\dots,\ b_s \end{array}\middle| q, z\right) = \sum_{n=0}^{\infty} \frac{(a_1,\dots,\ a_r;q)_n}{(q, b_1,\dots,\ b_s;q)_n} \left(-q^{(n-1)/2}\right)^{n(s+1-r)} z^n. \tag{3.10}$$

Finally, here is the result for entire functions in another parameter range:

Theorem 7. *For $r,\ s \in \mathbb{N}_0$, let $\alpha > 0,\ 1 > a_j,\ b_k > 0,\ 1 \le j \le r,\ 1 \le k \le s$ such that*

$$\prod_{j=1}^{r}(1-a_j)\prod_{k=1}^{s}(1-b_kq) \ge 4q^{2\alpha}, \tag{3.11}$$

then the entire function ${}_rA_s^{(\alpha)}(a_1,\dots a_r;\ b_1,\dots,\ b_s\ ;\ q;\ z)$ has all negative zeros.

4. Proofs

4.1. **Proof of Theorem 5.** For $n \in \mathbb{N}$, $0 < q < 1$ and $\alpha > 0$, since the sequences $\left\{(q^{-n};q)_k\,(-1)^k/(q;q)_k\right\}_{k=0}^{n}$ and $\left\{q^{\alpha k^2}/k!\right\}_{k=0}^{n}$ are PF sequences, then apply property 4 of Theorem 4 to get the PF sequence $\left\{(q^{-n};q)_k\,(-1)^k\,q^{\alpha k^2}/(q;q)_k\right\}_{k=0}^{n}$. Then by property 1 of Theorem 4 we know the polynomial

$$\sum_{k=0}^{n}\frac{(q^{-n};q)_k\,(-1)^k\,q^{\alpha k^2}z^k}{(q;q)_k} = A_q^{(\alpha)}\left(q^{-n};-z\right)$$

has all negative zeros, which is equivalent to that $A_q^{(\alpha)}\left(q^{-n};z\right)$ has all positive zeros.

For $m, \ell \in \mathbb{N},\ n_j \in \mathbb{N},\ 0 < q_j < 1,\ 1 \le j \le m$, and $\nu_r > -1,\ 0 < q_r < 1,\ 1 \le r \le \ell$, first by property 2, then by property 4 of Theorem 4 we know that

$$\left\{\prod_{j=1}^{m}\frac{\left(q_j^{-n_j};q_j\right)_k}{(q_j;q_j)_k}\frac{(-1)^{km}}{\prod_{r=1}^{\ell}\left(q_r,q_r^{\nu_r+1};q_r\right)_k}\right\}_{k\ge 0}$$

and

$$\left\{\prod_{j=1}^{m}\frac{\left(q_j^{-n_j};q_j\right)_k}{(q_j;q_j)_k}\frac{(-1)^{km}\,q^{\alpha k^2}}{\prod_{r=1}^{\ell}\left(q_r,q_r^{\nu_r+1};q_r\right)_k}\right\}_{k\ge 0}$$

are all PF sequences. Then for $\alpha \ge 0$ the polynomial (3.2) has all negative zeros.

For $m, \ell \in \mathbb{N}$, take

$$q_j = q,\ 1 \le j \le m, \quad q_r = q,\ 1 \le r \le \ell, \quad z = (1-q)^{2\ell}x,$$

and let $q \uparrow 1$ in

$$\sum_{k=0}^{\min\{n_j|1\leq j\leq m\}} \prod_{j=1}^{m}\prod_{s=1}^{k} \frac{\left(1-q^{s-1-n_j}\right)}{\left(1-q^{s}\right)} \frac{(-1)^{km}\, q^{\alpha k^2}\, (1-q)^{2k\ell}\, x^k}{\prod\limits_{r=1}^{\ell}\prod\limits_{t=1}^{k} (1-q^t)\,(1-q^{\nu_r+t})}$$

to get

$$\sum_{k=0}^{\min\{n_j|1\leq j\leq m\}} \prod_{j=1}^{m}\prod_{s=1}^{k} \frac{s-1-n_j}{s} \frac{(-1)^{km}\, x^k}{\prod\limits_{r=1}^{\ell}\prod\limits_{t=1}^{k} t(\nu_r+t)}$$

$$= \sum_{k=0}^{\min\{n_j|1\leq j\leq m\}} \frac{\prod_{j=1}^{m}(-n_j)_k}{\prod\limits_{r=1}^{\ell}(\nu_r+1)_k} \frac{(-1)^{km}\, x^k}{(k!)^{m+\ell}}$$

for each $z \in \mathbb{C}$. Then this limit is also uniform on any compact subset of $\mathbb{C}$ by Vitali's theorem 1. Thus the polynomial (3.3) has all negative zeros by applying Hurwitz's theorem 2.

4.2. **Proof of Theorem 6.** For $n \in \mathbb{N}$, $0 < q < 1$, $a \geq 0$ and $\alpha > 0$, since the sequences $\{(-a;q)_k / (q;q)_k\}_{k=0}^{n}$ and $\left\{q^{\alpha k^2}/k!\right\}_{k=0}^{n}$ are PF sequences, then apply property 4 of Theorem 4 to get the PF sequence $\left\{(-a;q)_k\, q^{\alpha k^2}/(q;q)_k\right\}_{k=0}^{n}$. Then by property 1 of Theorem 4 we know the polynomial

$$\sum_{k=0}^{n} \frac{(-a;q)_k\, q^{\alpha k^2}}{(q;q)_k} z^k$$

has all negative zeros. For $0 < q < 1$, $a \geq 0$ and $\alpha > 0$ and for each $n \in \mathbb{N}$ and $z \in \mathbb{C}$ we have,

$$\left|\sum_{k=0}^{n} \frac{(-a;q)_k\, q^{\alpha k^2}}{(q;q)_k} z^k\right| \leq \sum_{k=0}^{\infty} \frac{(-a;q)_k\, q^{\alpha k^2}}{(q;q)_k} |z|^k < \infty.$$

Now we first apply Vitali's theorem, then apply the Hurwitz's theorem to show the entire function $A_q^{(\alpha)}(-a;z)$ has no zeros outside the set $(-\infty, 0)$. Since by Lemma 14.1.4 of [11] we know that $A_q^{(\alpha)}(-a;z)$ has infinitely many zeros, then we have proved that $A_q^{(\alpha)}(-a;z)$ has infinitely many zeros and all of them are negative.

For $m, \ell \in \mathbb{N}_0$, $\alpha > 0$, $0 < q < 1$, $a_j \geq 0$, $0 < q_j < 1$, $1 \leq j \leq m$, and $\nu_r > -1$, $0 < q_r < 1$, $1 \leq r \leq \ell$, similar to the polynomial case, we first apply property 2, then apply property 4 of Theorem 4 we know that for each $n \in \mathbb{N}_0$, then sequence

$$\left\{\prod_{j=1}^{m} \frac{(-a_j;q_j)_k}{(q_j;q_j)_k} \frac{q^{\alpha k^2}}{\prod\limits_{r=1}^{\ell}\left(q_r, q_r^{\nu_r+1};q_r\right)_k}\right\}_{k=0}^{n}$$

is PF, thus the polynomial

$$\sum_{k=0}^{n}\prod_{j=1}^{m}\frac{(-a_j;q_j)_k}{(q_j;q_j)_k}\frac{q^{\alpha k^2}z^k}{\prod\limits_{r=1}^{\ell}\left(q_r,q_r^{\nu_r+1};q_r\right)_k}$$

has all negative zeros. For each $n\in\mathbb{N}_0$ and $z\in\mathbb{C}$ we have

$$\left|\sum_{k=0}^{n}\prod_{j=1}^{m}\frac{(-a_j;q_j)_k}{(q_j;q_j)_k}\frac{q^{\alpha k^2}z^k}{\prod\limits_{r=1}^{\ell}\left(q_r,q_r^{\nu_r+1};q_r\right)_k}\right|$$
$$\leq\sum_{k=0}^{\infty}\prod_{j=1}^{m}\frac{(-a_j;q_j)_k}{(q_j;q_j)_k}\frac{q^{\alpha k^2}|z|^k}{\prod\limits_{r=1}^{\ell}\left(q_r,q_r^{\nu_r+1};q_r\right)_k}<\infty.$$

Now we first apply Vitali's theorem, then apply Hurwitz's theorem to prove that the entire function (3.5) has no zeros outside $(-\infty,0)$. Since by Lemma 14.1.4 of [11] we know that the entire function (3.5) has infinitely many zeros, then the assertion for on (3.5) is proved.

Let

$$m\geq 0,\ \ell\geq 1,\ \alpha=\left(\ell+\frac{m}{2}\right),\ \nu_r\geq 0,\ 1\leq r\leq\ell,$$

and

$$a_j=0,\ q_j=q,\ 1\leq j\leq m,\quad q_r=q,\ 1\leq r\leq\ell$$

in (3.5), then for each $z\in\mathbb{N}$ and each $k\in\mathbb{N}_0$, by

$$\lim_{q\uparrow 1}\frac{(1-q)^k}{(q;q)_k}=\frac{1}{k!},\ \lim_{q\uparrow 1}\frac{(1-q)^k}{(q^{\nu_r+1};q)_k}=\frac{1}{(\nu_r+1)_k}$$

we have

$$\lim_{q\uparrow 1}\frac{q^{\alpha k^2}z^k(1-q)^{(m+2\ell)k}}{(q;q)_k^{m+\ell}\prod\limits_{r=1}^{\ell}(q^{\nu_r+1};q)_k}=\frac{z^k}{(k!)^{m+\ell}\prod\limits_{r=1}^{\ell}(\nu_r+1)_k}.$$

Since for $n\in\mathbb{N}$ we have

$$nq^{n-1}\leq\frac{1-q^n}{1-q}=1+q+\cdots+q^{n-1}\leq n,$$

and

$$n!q^{\binom{n}{2}}\frac{(q;q)_n}{(1-q)^n}\leq n!.$$

Then,

$$\frac{1}{n!}\leq\frac{(1-q)^n}{(q;q)_n}\leq\frac{q^{-\binom{n}{2}}}{n!}<\frac{q^{-\frac{n^2}{2}}}{n!}.$$

We also observe that for $\beta\geq 1$ we have

$$\frac{(1-q)^n}{(q^\beta;q)_n}\leq\frac{(1-q)^n}{(q;q)_n}.$$

Then,

$$\left|\frac{q^{\alpha k^2}z^k\,(1-q)^{(m+2\ell)k}}{(q;q)_k^{m+\ell}\prod_{r=1}^{\ell}\left(q^{\nu_r+1};q\right)_k}\right| \le \frac{|z|^k}{(k!)^{m+2\ell}}.$$

Hence

$$\lim_{q\uparrow 1}\sum_{k=0}^{\infty}\frac{q^{\alpha k^2}z^k\,(1-q)^{(m+2\ell)k}}{(q;q)_k^{m+\ell}\prod_{r=1}^{\ell}\left(q^{\nu_r+1};q\right)_k} = \sum_{k=0}^{\infty}\frac{z^k}{(k!)^{m+\ell}\prod_{r=1}^{\ell}(\nu_r+1)_k}$$

converges uniformly for z in any compact subset of $\mathbb{C}$. Then by Hurwitz's theorem we know that the entire function

$$g(z) = \sum_{k=0}^{\infty}\frac{z^k}{(k!)^{m+\ell}\prod_{r=1}^{\ell}(\nu_r+1)_k}$$

has no zeros outside $(-\infty, 0)$.

Let

$$a_k = \frac{1}{(k!)^{m+\ell}\prod_{r=1}^{\ell}(\nu_r+1)_k},$$

then by the Stirling's formula [1],

$$\log\Gamma(x+1) = x\log x + \mathcal{O}(x), \quad x\to\infty,$$

we have

$$-\log a_k = (m+2\ell)k\log k + \mathcal{O}(k)$$

as $k\to\infty$. Thus the order of $g(z)$ is given by [4]

$$\rho = \limsup_{k\to\infty}\frac{k\log k}{-\log|a_k|} = \frac{1}{m+2\ell},$$

which is a positive number less than $\frac{1}{2}$. Hence $g(z)$ has infinitely many zeros, and the claim on (3.6) is proved.

4.3. **Proof of Theorem 7.** By Lemma 14.1.4 of [11] we know that ${}_rA_s^{(\alpha)}(a_1,\dots a_r;\ b_1,\dots,\ b_s\ ;\ q;\ z)$ has infinitely many zeros.

Let

$$c_n = \frac{(a_1,\dots,\ a_r;\ q)_n}{(b_1,\dots,b_s;\ q)_n}q^{\alpha n^2},$$

then for $n\ge 1$ we have

$$\begin{aligned}\frac{c_n^2}{c_{n+1}c_{n-1}} &= q^{-2\alpha}\prod_{j=1}^{r}\frac{1-a_jq^{n-1}}{1-a_jq^n}\cdot\prod_{k=1}^{s}\frac{1-b_kq^n}{1-b_kq^{n-1}}\\ &\ge q^{-2\alpha}\prod_{j=1}^{r}(1-a_j)\prod_{k=1}^{s}(1-b_kq) \ge 4.\end{aligned}$$

Our assertion follows from Theorem B of [6] or Theorem A of [9].

References

[1] G. E. Andrews, R. Askey and R. Roy, *Special Functions,* Cambridge University Press, Cambridge, 1999.

[2] L. Ahlfors, Complex Analysis, 3rd edition, McGraw-Hill, 1979.

[3] M. Aissen, A. Edrei, I. J. Schoenberg, and A. M. Whitney, On the generating functions of totally positive sequences, Proc. Nat. Acad. Sci. U.S.A. 37 (1951), 303-307.

[4] R. P. Boas, Entire Functions, Academic Press, 1954.

[5] J. Carnicer, J. M. Peña and A. Pinkus, On some zero-increasing operators, Acta Math. Hungar. 94 (2002) 173-190.

[6] D. K. Dimitrov and J. M. Peña, Almost strict total positivity and a class of Hurwitz polynomials, Journal of Approximation Theory 132 (2005) 212-223.

[7] K. Driver, K. Jordaan and A. Martinínez-Finkelshtein, Pólya frequency sequences and real zeros of some ${}_3F_2$ polynomials, J. Math. Anal. 332(2007) 1045-1055.

[8] W. K. Hayman On the zeros of a q-Bessel function Complex Analysis and Dynamical Systems, Contemp. Math., Amer. Math. Soc., Providence, RI (2006), 205–216.

[9] J. I. Hutchinson, On a remarkable class of entire functions, Trans. Amer. Math. Soc. 25 (1923) 325–332.

[10] S. Karlin, Total Positioity, Vol. 1, Stanford University Press, 1968.

[11] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable,* Cambridge University Press, Cambridge, 2005.

[12] M. E. H. Ismail and R. Zhang, q-Bessel Functions and Rogers-Ramanujan Type Identities, Proceedings of American Mathematical Society, in production.

[13] E. H. Ismail and R. Zhang, q-Bessel Functions and Rogers-Ramanujan Type Identities, Proceedings of American Mathematical Society, in production.

[14] J. Pitman, Probabilistic bounds on the coefficients of polynomials with only real zeros, Journal of Combinatorial Theory Series. A 77 (1997) 279-303.

[15] E. C. Titchmarsh, The Theory of Functions, corrected second edition, Oxford University Press, Oxford, 1964.

Current address: College of Science, Northwest A&F University, Yangling, Shaanxi 712100, P. R. China.